

CLASS: 12	INDIAN SCHOOL MUSCAT FIRST PERIODIC ASSESSMENT	SUBJECT: MATHS
	SET - C	
QP.NO.	VALUE POINTS	SPLIT UP MARKS
1.	Let $x = \sec \theta \therefore \cot^{-1}(\cot \theta) = \theta = \sec^{-1} x$	(1 + 1) mks
2.	$f(x_1) = f(x_2) \Rightarrow 2x_1 = 2x_2 \Rightarrow x_1 = x_2 \Rightarrow f$ is one-one Let $y = 2x \Rightarrow x = \frac{y}{2} \notin \mathbb{N}$ (Eg: $y = 3$) $\Rightarrow \square f$ is not onto	1 mk 1 mk
3.	$gof = g(8x^3) = (8x^3)^{1/3} = 2x$ And $fog = f(x^{1/3}) = 8(x^{1/3})^3 = 8x$	1 mk 1 mk
4.	$\cos^{-1}\left(\cos\left(\pi + \frac{\pi}{6}\right)\right) = \cos^{-1}\left(-\cos\frac{\pi}{6}\right)$ $= \cos^{-1}\left(\cos\left(\pi - \frac{\pi}{6}\right)\right)$ $= \frac{5\pi}{6} \in [0, \pi]$	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2}$ mk $\frac{1}{2}$ mk
5.	$\tan^{-1}\left(\frac{\frac{x-1}{x-2} + \frac{x+1}{x+2}}{1 - \left(\frac{x-1}{x-2}\right)\left(\frac{x+1}{x+2}\right)}\right) = \frac{\pi}{4}, \quad \frac{x^2-1}{x^2-4} < 1$ Solving to get $2x^2 - 4 = -3$ Final answer : $x = \pm \frac{1}{\sqrt{2}}$	2 mks 1 mk 1 mk
6.	Let $y = 9x^2 + 6x - 5 = (3x + 1)^2 - 6 \Rightarrow x = \frac{\sqrt{y+6}-1}{3}$ Proving $fog = y$ and $gof = x$ $\Rightarrow f$ is invertible $\therefore f^{-1} = g = \frac{\sqrt{y+6}-1}{3}$	1 mk (1+1) mks $\frac{1}{2}$ mk $\frac{1}{2}$ mk
7.	For proving Reflexive For proving Symmetric For proving Transitive Conclusion	1 mk 1 mk 1 $\frac{1}{2}$ mk $\frac{1}{2}$ mk