

SET	C
-----	---

INDIAN SCHOOL MUSCAT  
HALF YEARLY EXAMINATION 2023  
MATHEMATICS - 041

CLASS:X

Max.Marks: 80

MARKING SCHEME			
SET	QN.NO	VALUE POINTS	MARKS SPLIT UP
	1	(c) $60^\circ$	
	2	(c) $2 \times 7^2$	
	3	(a) 240	
	4	(d) 10	
	5	(c) $\frac{12}{13}$	
	6	(b) AA similarity criterion	
	7	(a) $+2\sqrt{3}, -2\sqrt{3}$	
	8	(c) 20	
	9	(a) $3:1$	
	10	(b) -1	
	11	(d) 4 units	
	12	(a) 16 cm	
	13	(b) 6	
	14	(a) Both assertion (A) and reason (R) are true and reason (R) is the correct explanation of assertion (A).	
	15	(d) $\frac{1}{7}$	
	16	(b) 10 cm	

	17	(b) 2	
	18	(a) 26	
	19	(d) Assertion (A) is false but reason(R) is true.	
	20	(b) 1.5	
	21	$S = (3 + \sqrt{2}) + (3 - \sqrt{2}) = 6$ $P = (3 + \sqrt{2}) \times (3 - \sqrt{2}) = (3)^2 - (\sqrt{2})^2 = 9 - 2 = 7$ Quadratic polynomial = $x^2 - Sx + P = x^2 - 6x + 7$	$\frac{1}{2}$ $\frac{1}{2}$ 1
	22	AB = 10 units ... [Given $AB^2 = 10^2 = 100$ ... [Squaring both sides $(11 - 3)^2 + (y + 1)^2 = 100$ $8^2 + (y + 1)^2 = 100$ $(y + 1)^2 = 100 - 64 = 36$ $y + 1 = \pm 6$ ... [Taking square-root on both sides $y = -1 \pm 6 \therefore y = -7 \text{ or } 5$ OR Area of $\triangle ABC = \frac{1}{2} \times \text{base} \times \text{corr, altitude}$ $= \frac{1}{2} \times 5 \times 3 = 7.5 \text{ sq. units}$	1     1  1  1
	23	Table Median = 340	1 1
	24	HCF = 10 LCM = 300	1 1
	25	$a = 2, b = -4, c = 4$ $b^2 - 4ac = -16 < 0$ No real root OR Roots are $\frac{2}{3}$ and $-\frac{1}{2}$	$\frac{1}{2}$ 1 $\frac{1}{2}$
	26	Volume of cone Volume of cylinder Volume of hemisphere Total volume Conclusion	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 1 $\frac{1}{2}$

27	<p><b>Given:</b> ABCD is parallelogram circumscribing a circle.</p> <p><b>To prove:</b> ABCD is a rhombus</p> <p><b>Proof:</b> We have, <math>DR = DS</math> ... (i)</p> <p>[Lengths of tangents drawn from an external point to a circle are equal]</p> <p>Also, <math>AP = AS</math> ... (ii)</p> <p><math>BP = BQ</math> ... (iii)</p> <p><math>CR = CQ</math> ... (iv)</p> <p>Adding (i), (ii), (iii) and (iv),</p> $(DR + CR) + (AP + BP) = (DS + AS) + (BQ + CQ)$ $\Rightarrow CD + AB = AD + BC$ $\Rightarrow 2AB = 2AD \quad [\because \text{In parallelogram, opposite sides are equal}]$ $\Rightarrow AB = AD$ <p><math>\therefore AB = AD = BC = CD</math></p> <p>Hence, ABCD is a rhombus as all sides are equal in rhombus.</p> <p style="text-align: center;">OR</p> <p><b>Given:</b> A quadrilateral ABCD which circumscribes a circle.</p> <p>Let it touches the circle at P, Q, R and S as shown in figure.</p> <p><b>To Prove:</b> <math>AB + CD = AD + BC</math></p> <p><b>Proof:</b> We know that the lengths of the tangents drawn from a point outside the circle to the circle are equal.</p> <p><math>\therefore AP = AS; BP = BQ; CQ = CR</math> and <math>DR = DS</math> ... (i)</p> <p>Consider, <math>AB + CD = AP + PB + CR + RD</math></p> $= AS + BQ + CQ + DS$ <p>[using (i)]</p> $= (AS + DS) + (BQ + CQ) = AD + BC$	<p>Fig 1/2</p> <p>1</p> <p>1/2</p> <p>1/2</p>
28	<p>We have, <math>6x^2 - 3 - 7x</math></p> $= 6x^2 - 7x - 3$ $= (2x - 3)(3x + 1)$ <p>Zeroes are:</p> $2x - 3 = 0 \text{ or } 3x + 1 = 0$ <p>Therefore <math>x = 3/2</math> or <math>x = -1/3</math></p> <p>Verification:</p> <p>Here <math>a = 6, b = -7, c = -3</math></p> <p>Sum of the zeroes, <math>(\alpha + \beta) = 3/2 + (-1/3) = (9 - 2)/6 = 7/6</math></p> $7/6 = -(\text{coefficient of } x) / (\text{Coefficient of } x^2) = -b/a$ <p>Product of Zeroes <math>(\alpha \times \beta) = 3/2 \times (-1/3) = -3/6</math></p> $-3/6 = \text{Constant term} / \text{Coefficient of } x^2 = c/a$ <p>Therefore, Relationship holds</p>	<p>Fig 1/2</p> <p>1</p> <p>1</p> <p>1</p>



34

Class interval	Mid-values ( $x_i$ )	Frequency ( $f_i$ )	$u_i = \frac{x_i - 50}{20}$	$f_i u_i$
0-20	10	17	-2	-34
20-40	30	$f_1$	-1	$-f_1$
40-60	50	32	0	0
60-80	70	$f_2$	1	$f_2$
80-100	90	19	2	38
<b>Total</b>		$\Sigma f_i = 68 + f_1 + f_2$		$\Sigma f_i u_i = 4 - f_1 + f_2$

$$f_1 + f_2 = 52 \text{ ----(i)}$$

$$\text{Mean} = 50$$

$$\text{Therefore, } f_1 - f_2 = 4 \text{ -----(ii)}$$

Solving we get  $f_1 = 28$  and  $f_2 = 24$

OR

Index	No. of weeks ( $f_i$ )	c.f.
1500-1600	3	3
1600-1700	11 $f_0$	14
1700-1800	12 $f_1$	26
1800-1900	7 $f_2$	33
1900-2000	9	42
2000-2100	8	50
2100-2200	2	52
	$\Sigma f_i = 52$	

$$n = 52, \frac{n}{2} = \frac{52}{2} = 26$$

$\therefore$  Median class is 1700-1800

$$\begin{aligned} \therefore \text{Median} &= l + \frac{\frac{n}{2} - c.f.}{f} \times h \\ &= 1700 + \left( \frac{12}{12} \times 100 \right) = \mathbf{1800} \end{aligned}$$

$\therefore$  Maximum frequency is 12

$\Rightarrow$  Modal class is 1700-1800

$$\begin{aligned} \text{Mode} &= l + \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \times h \\ &= 1700 + \frac{12 - 11}{24 - 11 - 7} \times 100 \end{aligned}$$

$$= \mathbf{1716.\bar{6} \text{ or } 1716.67 \text{ (approx.)}}$$



	<b>36</b>	(i)50m (ii)30m (iii) 24m OR (iii)36m	1 1 2
	<b>37</b>	(i)0 units (ii)(4,2) (iii)Ramesh travels more OR (iii) Library	1 1 2
	<b>38</b>	(i)distance = (speed )x time (ii) $x^2 + 30x - 400 = 0$ (iii)10 km/hour OR (iii)1.5 hour	1 1 2